



## Article 134

Technical Note #20 from *Watershed Protection Techniques*. 1(2): 73-75

# Minimizing the Impact of Golf Courses on Streams

Over 13,000 golf courses now exist in the U.S. and many more will be constructed to meet the growing popularity of the sport. The construction of a new golf course has the potential to create adverse impacts on the aquatic environment. To begin with, a typical 18-hole golf course can convert as much as 100 acres of rural land into a highly “terra-formed” environment of fairways, greens, tees, sand traps, and water hazards. As such, golf courses are often an attractive part of the urban landscape. Haphazardly designed golf courses, however, can disrupt and degrade the wetlands, floodplains, riparian zones, and forests that contribute to stream quality.

A second recurring concern about golf courses are the large inputs of fertilizer, pesticides, fungicides, and other chemicals that are required to maintain vigorous and attractive greens. In many cases, chemical application rates can rival and even exceed those used in intensive agriculture. Table 1 shows a side by side comparison of chemical application rates for a coastal plain golf course and cropland in Maryland, as reported by Klein (1990).

The actual rate of fertilizer and pesticide application rates at a particular golf course can vary considerably, depending on the soil, climate, and management program. As an example, fungicides and nematicides are only lightly used in regions with cold winters, but constitute a major fraction of total pesticide applications in warmer climates. Given such intensive use of chemicals, golf courses clearly have the potential to deliver pollutants to ground and surface waters. Actual monitoring data on pollutant loads from golf courses, however, are quite scarce.

Golf courses are also intensive water consumers, particularly in drier regions of the country. This need for irrigable water can place strong demands on local groundwater and/or surface water supplies, which in turn, can cause baseflow depletion. In addition, the construction of the ubiquitous golf course water hazards can lead to downstream warming in sensitive trout streams.

In the late 1980s, Baltimore County, Maryland was confronted with a wave of golf course development proposals and strong concerns about the possible risk they might have on their Piedmont streams. The Department of Environmental Protection and Resource Management drafted and revised a series of environmental guidelines for new golf course construction. The guidelines stress the importance of integrating the layout of the course with the natural features of the site.

For example, the guidelines require a detailed evaluation of wetlands, perennial and intermittent streams, floodplains, slopes, forest stands and habitat features at the proposed course. The course must be configured to avoid or minimize disturbance to these resource areas. In this respect, long broad fairways are a prime culprit, as they frequently cross or encroach into streams and other buffer areas.

Consequently, the guidelines devote a great deal of attention to the issue of fairway crossings (see Figure 1). For example, no more than two fairway crossings are allowed for each 1,000 feet of stream length. These crossing must be perpendicular to the stream. If forests or wetlands are present at the crossing, this zone must be managed as unplayable rough and remain undis-

**Table 1: Comparative Chemical Application Rates for a Maryland Golf Course and Corn/Soybean Rotation Reported in Pounds/Acre/Year (Klein, 1990)**

Chemical	Cropland	Fairway	Greens	Tees
Nitrogen	184	150	213	153
Phosphorus	80	88	44	93
Herbicides	5.8	10.4	10.2	11.4
Insecticide	1.0	2.0	2.0	2.0
Fungicide	0.0	26.9	34.9	26.9
Total Pesticides	5.8	37.3	45.1	38.3